

Chlamydia Screening in Women (CHL)

Measure Description

The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

Numerator Compliance

At least one chlamydia test during the measurement year.

Best Practices

- ★ **Incorporate** universal screening of all members in this age range.
- ★ **Screen** should occur with or without symptoms.
- ★ **Screen** if oral contraceptives, sexually transmitted diseases (STD)/ sexually transmitted infections (STI) or urinary symptoms are discussed.
- ★ **Use** phone, email, or other means to have patients come in for their routine yearly visit.



Quality Value Set Directory Coding Tips

Chlamydia Tests

CPT

87110, 87270, 87320, 87490, 87491, 87492, 87810

LOINC

14463-4, 14464-2, 14465-9, 14467-5, 14474-1, 14513-6, 16600-9, 21190-4, 21191-2, 21613-5, 23838-6, 31775-0, 34710-4, 42931-6, 43304-5, 43404-3, 44806-8, 44807-6, 45068-4, 45069-2, 45072-6, 45073-4, 45075-9, 45084-1, 45089-0, 45090-8, 45091-6, 45093-2, 45095-7, 4993-2, 50387-0, 53925-4, 53926-2, 57287-5, 6353-7, 6356-0, 6357-8, 80360-1, 80361-9, 80362-7, 80363-5, 80364-3, 80365-0, 80367-6, 82306-2, 87949-4, 87950-2, 88221-7, 89648-0, 91860-7, 91873-0

Members recommended for routine chlamydia screening

Female

AdministrativeGender

Female

If used: both required

Sex assigned at birth

LOINC

76689-9

Female

LOINC

LA3-6

Female sex for clinical use

SexParameterForClinicalUse

Female-typical