

Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)

Measure Description

The percentage of members 18–64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

Numerator Compliance

A glucose test or an HbA1c test performed during the measurement year.

Best Practices

- ★ **Order** a diabetes screening test every year and build care gap “alerts” in your electronic medical record.
- ★ **Follow up** with patients to discuss and educate on lab results.
- ★ **Coordinate** care with patients’ other providers.
- ★ **Outreach** patients who cancel appointments and reschedule them as soon as possible.
- ★ **Encourage** shared decision-making by educating patients and caregivers about increased risk of diabetes with antipsychotic medications.
- ★ **Discuss** with patient the importance of screening for diabetes.
- ★ **Educate** the patient on symptoms of new onset diabetes.



Quality Value Set Directory Coding Tips

Glucose Lab Test

CPT

80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

LOINC

10450-5, 1492-8, 1494-4, 1496-9, 1499-3, 1501-6, 1504-0, 1507-3, 1514-9, 1518-0, 1530-5, 1533-9, 1554-5, 1557-8, 1558-6, 17865-7, 20436-2, 20437-0, 20438-8, 20440-4, 2345-7, 26554-6, 41024-1, 49134-0, 6749-6, 9375-7

HbA1c Lab Test

CPT

83036, 83037

LOINC

17855-8, 17856-6, 4548-4, 4549-2, 96595-4

HbA1c Test Result or Finding

CPT-CAT-II

3044F, 3046F, 3051F, 3052F