

Syphilis Screening of Sexually Active Patients (SYP)

Measure Description

The percentage of patients 18–64 years of age who were identified as sexually active and who had at least one syphilis test during the measurement year.

Numerator Compliance

At least one syphilis test during the measurement year.

Best Practices

- ★ **Utilize** annual wellness visits to conduct a complete sexual history assessment and order test.
- ★ **Screenings** should also occur at any visit where oral contraceptives, sexually transmitted diseases (STD)/sexually transmitted infections (STI) or urinary symptoms are discussed.
- ★ **Use** phone, email, or other means to have patients come in for their routine yearly visit.
- ★ **Encourage** the member to make an appointment for STI screening once a year (or, to keep their appointment if they indicate they already have one).
- ★ **Reschedule** no shows immediately.



Quality Value Set Directory Coding Tips

Syphilis test, non-treponemal antibody; qualitative
CPT
 86592

Syphilis test, non-treponemal antibody; quantitative
CPT
 86593

Treponema pallidum
CPT
 86780

Reagin Ab Titr Ser
LOINC
 11084-1

RPR Ser QI
LOINC
 20507-0

RPR Ser-Titr
LOINC
 31147-2

T pallidum Ab Ser LA-aCnc
LOINC
 17725-3

T pallidum Ab Ser QI HA
LOINC
 8041-6

T pallidum Ab Ser QI IA
LOINC
 24110-9

T pallidum IgG Ser QI IA
LOINC
 47238-1

T pallidum IgG+IgM Ser QI
LOINC
 34147-9

T pallidum IgG+IgM Ser QI IA
LOINC
 47236-5