

Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)

Measure Description

The percentage of patients 18–64 years of age with schizophrenia or schizoaffective disorder and diabetes who had both an LDL-C test and an HbA1c test during the measurement year.

Numerator Compliance

An HbA1c test and an LDL-C test were performed during the measurement year. The patient must have both tests.

Best Practices

- ★ **Order** a diabetes screening test every year and build care gap “alerts” in your electronic medical record.
- ★ **Follow up** with patients to discuss and educate on lab results.
- ★ **Coordinate** care with patients’ other providers.
- ★ **Outreach** patients who cancel appointments and reschedule them as soon as possible.
- ★ **Encourage** shared decision-making by educating patients and caregivers about increased risk of diabetes with antipsychotic medications.
- ★ **Discuss** with patient the importance of screening for diabetes.
- ★ **Educate** the patient on symptoms of new onset diabetes.



Quality Value Set Directory Coding Tips

HbA1c Lab Test

LOINC

17855-8, 17856-6, 4548-4, 4549-2, 96595-4

CPT

83036, 83037

HbA1c Test Result or Finding

CPT-CAT-II

3044F, 3046F, 3051F, 3052F

LDL-C Lab Test

CPT

80061, 83700, 83701, 83704, 83721

LOINC

12773-8, 13457-7, 18261-8, 18262-6, 2089-1, 49132-4, 55440-2, 96259-7

LDL-C Test Result or Finding

CPT-CAT-II

3048F, 3049F, 3050F, 3051F, 3052F