

Chlamydia Screening (CHL)

Measure Description

The percentage of members 16–45 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

Numerator Compliance

At least one chlamydia test during the measurement year.



Quality Value Set Directory Coding Tips

Chlamydia Tests

CPT

87110, 87270, 87320, 87490, 87491, 87492, 87810

LOINC

14463-4, 14464-2, 14465-9, 14467-5, 14474-1, 14513-6, 16600-9, 21190-4, 21191-2, 21613-5, 23838-6, 31775-0, 34710-4, 42931-6, 43304-5, 43404-3, 44806-8, 44807-6, 45068-4, 45069-2, 45072-6, 45073-4, 45075-9, 45084-1, 45089-0, 45090-8, 45091-6, 45093-2, 45095-7, 4993-2, 50387-0, 53925-4, 53926-2, 57287-5, 6353-7, 6356-0, 6357-8, 80360-1, 80361-9, 80362-7, 80363-5, 80364-3, 80365-0, 80367-6, 82306-2, 87949-4, 87950-2, 88221-7, 89648-0, 91860-7, 91873-0

Chlamydia Screening

Administrative Gender

Female

LOINC

76689-9

NYS Chlamydia Screen

CPT

87110, 87270, 87320, 87490, 87491, 87492, 87810, 86631, 86332, 87800, 3511F, 0353U

Best Practices

- ★ **Incorporate** universal screening of all members in this age range.
- ★ **Screening** should occur with or without symptoms.
- ★ **Screening** should also occur at any visit where oral contraceptives, sexually transmitted diseases (STD)/ sexually transmitted infections (STI) or urinary symptoms are discussed.
- ★ **Use** phone, email, or other means to have members come in for their routine yearly visit.