

The following diagnosis and/or procedure codes in the HEDIS Quick Reference Guide are in compliance with the HEDIS® 2017 Volume 2 Technical Specifications. ICD-10 codes must be used effective October 1, 2015. ICD-9 codes are included for certain measures because they remain in effect for 2017 HEDIS.

### ANTIDEPRESSANT MEDICATION MANAGEMENT (AMM)

The percentage of members 18 years of age and older with a diagnosis of major depression and were treated with antidepressant medication, and who remained on an antidepressant medication treatment. Two rates are reported.

1. Effective Acute Phase Treatment. The percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks).
2. Effective Continuation Phase Treatment. The percentage of members who remained on an antidepressant medication for at least 180 days (6 months).

Code set	
ICD-9-CM Diagnosis	296.20-296.25, 296.30-296.35, 298.0, 311
ICD-10-CM Diagnosis	F32.0-F32.4, F32.9, F33.0-F33.3, F33.41, F33.9
CPT	
AMM Stand Alone Visits: 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99384-99387, 99394-99397, 99401-99404, 99411, 99412, 99510 ED Visits: 99281-99285	AMM Stand Alone Visits: G0155, G0176, G0177, G0409-G0411, G0463, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, M0064, S0201, S9480, S9484, S9485, T1015
CPT	
AMM Visits: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90867-90870, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255	<i>WITH</i>
POS	
AMM Visits: 03, 05, 07, 09, 11-20, 22, 24, 33, 49, 50, 52, 53, 71, 72	
Antidepressant Medications: The following list of medications is based upon the 2017 HEDIS® Technical Specifications.	
Description	Prescription
Miscellaneous Antidepressants	<ul style="list-style-type: none"> <li>• Bupropion</li> <li>• Vilazodone</li> <li>• Vortioxetine</li> </ul>
Monoamine Oxidase Inhibitors	<ul style="list-style-type: none"> <li>• Isocarboxazid</li> <li>• Selegiline</li> <li>• Phenelzine</li> <li>• Tranylcypromine</li> </ul>
Phenylpiperazine Antidepressants	<ul style="list-style-type: none"> <li>• Nefazodone</li> <li>• Trazodone</li> </ul>
Psychotherapeutic combinations	<ul style="list-style-type: none"> <li>• Amitriptyline-chlordiazepoxide</li> <li>• Amitriptyline-perphenazine</li> <li>• Fluoxetine-olanzapine</li> </ul>
SNRI Antidepressants	<ul style="list-style-type: none"> <li>• Desvenlafaxine</li> <li>• Levomilnacipran</li> <li>• Duloxetine</li> <li>• Venlafaxine</li> </ul>
SSRI antidepressants	<ul style="list-style-type: none"> <li>• Citalopram</li> <li>• Escitalopram</li> <li>• Fluoxetine</li> <li>• Fluvoxamine</li> <li>• Paroxetine</li> <li>• Sertraline</li> </ul>
Tetracyclic antidepressants	<ul style="list-style-type: none"> <li>• Maprotiline</li> <li>• Mirtazapine</li> </ul>
Tricyclic antidepressants	<ul style="list-style-type: none"> <li>• Amitriptyline</li> <li>• Amoxapine</li> <li>• Clomipramine</li> <li>• Desipramine</li> <li>• Doxepin (&gt;6mg)</li> <li>• Imipramine</li> <li>• Nortriptyline</li> <li>• Protriptyline</li> <li>• Trimipramine</li> </ul>

## FOLLOW-UP AFTER HOSPITALIZATION FOR MENTAL ILLNESS (FUH)

<p><i>The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner. Two rates are reported:</i></p> <ul style="list-style-type: none"> <li>• The percentage of discharges for which the member received follow-up within 7 days of discharge.</li> <li>• The percentage of discharges for which the member received follow-up within 30 days of discharge.</li> </ul>		
Code set		
Mental Health Practitioner	<ul style="list-style-type: none"> <li>• An MD or doctor of osteopathy (DO) who is certified as a psychiatrist or child psychiatrist by the American Medical Specialties Board of Psychiatry and Neurology or by the American Osteopathic Board of Neurology and Psychiatry; or, if not certified, who successfully completed an accredited program of graduate medical or osteopathic education in psychiatry or child psychiatry and is licensed to practice patient care psychiatry or child psychiatry, if required by the state of practice.</li> <li>• An individual who is licensed as a psychologist in his/her state of practice, if required by the state of practice.</li> <li>• An individual who is certified in clinical social work by the American Board of Examiners; who is listed on the National Association of Social Worker's Clinical Register; or who has a master's degree in social work and is licensed or certified to practice as a social worker, if required by the state of practice.</li> <li>• A registered nurse (RN) who is certified by the American Nurses Credentialing Center (a subsidiary of the American Nurses Association) as a psychiatric nurse or mental health clinical nurse specialist, or who has a master's degree in nursing with a specialization in psychiatric/mental health and two years of supervised clinical experience and is licensed to practice as a psychiatric or mental health nurse, if required by the state of practice.</li> <li>• An individual (normally with a master's or a doctoral degree in marital and family therapy and at least two years of supervised clinical experience) who is practicing as a marital and family therapist and is licensed or a certified counselor by the state of practice, or if licensure or certification is not required by the state of practice, who is eligible for clinical membership in the American Association for Marriage and Family Therapy.</li> <li>• An individual (normally with a master's or doctoral degree in counseling and at least two years of supervised clinical experience) who is practicing as a professional counselor and who is licensed or certified to do so by the state of practice, or if licensure or certification is not required by the state of practice, is a National Certified Counselor with a Specialty Certification in Clinical Mental Health Counseling from the National Board for Certified Counselors (NBCC).</li> </ul>	
ICD-9-CM Diagnosis	295.00-299.91, 300.3, 300.4, 301.0, 301.10-301.13, 301.20-301.22, 301.3, 301.4, 301.50-301.59, 301.6, 301.7, 301.81-301.89, 301.9, 308.0-308.9, 309.0, 309.1, 309.21-309.29, 309.3, 309.4, 309.81-309.89, 309.9, 311, 312.00-314.9	
ICD-10-CM Diagnosis	Use the appropriate code family: F	
CPT		HCPCS
Follow-up visits identified by the following CPT or HCPCS codes must be with a mental health practitioner within 7 and 30 days of discharge.		
Stand Alone Visits: 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99383-99387, 99393-99397, 99401-99404, 99411, 99412, 99510 Transitional Care: 99495 (14 Days, used for 30-Day Follow-up indicator) and 99496 (7 Days)	G0155, G0176, G0177, G0409-G0411, G0463, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, M0064, S0201, S9480, S9484, S9485, T1015	
CPT		POS
Follow-up visits identified by the following CPT/POS codes must be with a mental health practitioner.		
90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90867-90870, 90875, 90876	WITH	03, 05, 07, 09, 11-20, 22, 24, 33, 49, 50, 52, 53, 71, 72
99221-99223, 99231-99233, 99238, 99239, 99251-99255	WITH	52, 53

## INITIATION AND ENGAGEMENT OF ALCOHOL AND OTHER DRUG DEPENDENCE TREATMENT (IET)

<p><i>The percentage of adolescent and adult members (ages 13 and older) with a new episode of alcohol or other drug (AOD) dependence who received the following:</i></p> <ul style="list-style-type: none"> <li>• <i>Initiation of AOD Treatment. The percentage of members who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis.</i></li> <li>• <i>Engagement of AOD Treatment. The percentage of members who initiated treatment and who had two or more additional services with a diagnosis of AOD within 30 days of the initiation visit.</i></li> </ul>		
Code set		
ICD-9-CM Diagnosis	AOD: 291.0-291.5, 291.81-291.89, 291.9, 303.00-303.02, 303.90, 303.91, 303.92, 304.00-304.02, 304.10-304.12, 304.20-304.22, 304.30-304.32, 304.40-304.42, 304.50-304.52, 304.60-304.62, 304.70-304.72, 304.80-304.82, 304.90-304.92, 305.00-305.02, 305.20-305.22, 305.30-305.32, 305.40-305.42, 305.50-305.52, 305.60-305.62, 305.70-305.72, 305.80-305.82, 305.90-305.92, 535.30, 535.31, 571.1	
ICD-10-CM Diagnosis	Use the appropriate code family: F	

CPT		HCPCS
Stand Alone Visits identified by the following CPT or HCPCS <i>combined with</i> AOD Diagnosis Code		
98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99384-99387, 99394-99397, 99401-99404, 99408, 99409, 99411, 99412, 99510	G0155, G0176, G0177, G0396, G0397, G0409-G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0020, H0022, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9485, T1006, T1012, T1015	
CPT		POS
IET visits identified by the following CPT/POS codes <i>combined with</i> AOD Diagnosis Code		
90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876	WITH	03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 57, 71, 72
99221-99223, 99231-99233, 99238, 99239, 99251-99255	WITH	52, 53

### ADHERENCE TO ANTIPSYCHOTIC MEDICATIONS FOR INDIVIDUALS WITH SCHIZOPHRENIA (SAA)

The percentage of members ages 19-64 during the measurement year with schizophrenia who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.		
ICD-9-CM Diagnosis	295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95	
ICD-10-CM Diagnosis	F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9	
HCPCS	14 day supply: J2794 28 day supply: J0401, J1631, J2358, J2426, J2680	
Antipsychotic Medications:		
Description	Prescription	Days Supply
Miscellaneous antipsychotic agents (oral)	<ul style="list-style-type: none"> <li>• Aripiprazole</li> <li>• Asenapine</li> <li>• Brexpiprazole</li> <li>• Cariprazine</li> <li>• Clozapine</li> <li>• Haloperidol</li> <li>• Iloperidone</li> <li>• Loxapine</li> <li>• Lurasidone</li> <li>• Molindone</li> <li>• Olanzapine</li> <li>• Paliperidone</li> <li>• Pimozide</li> <li>• Quetiapine</li> <li>• Quetiapine fumarate</li> <li>• Risperidone</li> <li>• Ziprasidone</li> </ul>	
Phenothiazine antipsychotics (oral)	<ul style="list-style-type: none"> <li>• Chlorpromazine</li> <li>• Fluphenazine</li> <li>• Perphenazine</li> <li>• Perphenazine-amitriptyline</li> <li>• Prochlorperazine</li> <li>• Thioridazine</li> <li>• Trifluoperazine</li> </ul>	
Psychotherapeutic combinations (oral)	<ul style="list-style-type: none"> <li>• Fluoxetine-olanzapine</li> </ul>	
Thioxanthenes (oral)	<ul style="list-style-type: none"> <li>• Thiothixene</li> </ul>	
Long-acting injections	<ul style="list-style-type: none"> <li>• Aripiprazole</li> <li>• Fluphenazine decanoate</li> <li>• Haloperidol decanoate</li> <li>• Olanzapine</li> <li>• Paliperidone palmitate</li> </ul>	28 days supply
	<ul style="list-style-type: none"> <li>• Risperidone</li> </ul>	14 days supply
Exclusions include members who met at least one of the following during the measurement year: 1) <i>Dementia diagnosis.</i> 2) <i>Did not receive at least two (2) antipsychotic medication dispensing events.</i>		
ICD-9-CM Diagnosis	290.0, 290.10-290.13, 290.20, 290.21, 290.3, 290.40-290.43, 290.8, 290.9, 291.2, 292.82, 294.0, 294.10, 294.11, 294.20, 294.21, 331.0, 331.82	
ICD-10-CM Diagnosis	Use the appropriate code family: F, G	

## DIABETES SCREENING FOR PEOPLE WITH SCHIZOPHRENIA OR BIPOLAR DISORDER WHO ARE USING ANTIPSYCHOTIC MEDICATIONS (SSD)

<i>Percentage of members ages 18-64 with schizophrenia or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test.</i>	
ICD-9-CM Diagnosis	Schizophrenia: 295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95 Bipolar Disorder: 296.00-296.06, 296.10-296.16, 296.40-296.46, 296.50-296.56, 296.60-296.66, 296.7 Other Bipolar Disorders: 296.80-296.82, 296.89
ICD-10-CM Diagnosis	Schizophrenia: F20.0-F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Bipolar Disorder: F30.10-F30.13, F30.2-F30.4, F30.8, F30.9, F31.0, F31.10-F31.13, F31.2, F31.30-F31.32, F31.4, F31.5, F31.60-F31.64, F31.70-F31.78 Other Bipolar Disorders: F31.81, F31.89, F31.9
CPT	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951, 83036, 83037
CPT II	3044F, 3045F, 3046F
<b>Antipsychotic Medications:</b>	
Miscellaneous antipsychotic agents	<ul style="list-style-type: none"> <li>• Aripiprazole</li> <li>• Asenapine</li> <li>• Brexpiprazole</li> <li>• Cariprazine</li> <li>• Clozapine</li> <li>• Haloperidol</li> <li>• Iloperidone</li> <li>• Loxapine</li> <li>• Lurasidone</li> <li>• Molindone</li> <li>• Olanzapine</li> <li>• Paliperidone</li> <li>• Pimozide</li> <li>• Quetiapine</li> <li>• Quetiapine fumarate</li> <li>• Risperidone</li> <li>• Ziprasidone</li> </ul>
Phenothiazine antipsychotics	<ul style="list-style-type: none"> <li>• Chlorpromazine</li> <li>• Fluphenazine</li> <li>• Perphenazine</li> <li>• Perphenazine-amitriptyline</li> <li>• Prochlorperazine</li> <li>• Thioridazine</li> <li>• Trifluoperazine</li> </ul>
Psychotherapeutic combinations	<ul style="list-style-type: none"> <li>• Fluoxetine-Olanzapine</li> </ul>
Thioxanthenes	<ul style="list-style-type: none"> <li>• Thiothixene</li> </ul>
Long-acting injections	<ul style="list-style-type: none"> <li>• Aripiprazole</li> <li>• Fluphenazine decanoate</li> <li>• Haloperidol decanoate</li> <li>• Olanzapine</li> <li>• Paliperidone palmitate</li> <li>• Risperidone</li> </ul>

## DIABETES MONITORING FOR PEOPLE WITH DIABETES AND SCHIZOPHRENIA (SMD)

<i>The percentage of members ages 18-64 with schizophrenia and diabetes who had both an LDL-C and an HbA1c test during the measurement year.</i>	
ICD-9-CM Diagnosis	Schizophrenia: 295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95 Diabetes: 250.00-250.03, 250.10-250.13, 250.20-250.23, 250.30-250.33, 250.40-250.43, 250.50-250.53, 250.60-250.63, 250.70-250.73, 250.80-250.83, 250.90-250.93, 357.2, 362.01-362.07, 366.41, 648.00-648.04
ICD-10-CM Diagnosis	Schizophrenia: F20.0-F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Diabetes: Use the appropriate code family: E or O (pre-existing DM in pregnancy)
CPT	HbA1c tests: 83036, 83037 LDL-C tests: 80061, 83700, 83701, 83704, 83721
CPT II	3044F, 3045F, 3046F, 3048F, 3049F, 3050F

## CARDIOVASCULAR MONITORING FOR PEOPLE WITH CARDIOVASCULAR DISEASE AND SCHIZOPHRENIA (SMC)

<i>The percentage of members 18–64 years of age with schizophrenia and cardiovascular disease, who had an LDL-C test during the measurement year.</i>	
ICD-9-CM Diagnosis	Schizophrenia: 295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95  Ischemic Vascular Disease (IVD): 411.0, 411.1, 411.81, 411.89, 413.0, 413.9, 414.00-414.07, 414.2-414.4, 414.8, 414.9, 429.2, 433.00, 433.01, 433.10, 433.11, 433.20, 433.21, 433.30, 433.31, 433.80, 433.81, 433.90, 433.91, 434.00, 434.01, 434.10, 434.11, 434.90, 434.91, 437.0, 440.0, 440.1, 440.20-440.24, 440.29-440.32, 440.4, 440.8, 440.9, 444.01, 444.09, 444.1, 444.21, 444.22, 444.81, 444.89, 444.9, 445.01, 445.02, 445.81, 445.89
ICD-10-CM Diagnosis	Schizophrenia: F20.0-F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Ischemic Vascular Disease (IVD): Use the appropriate code family: I
CPT Code Set	
LDL-C Test	80061, 83700, 83701, 83704, 83721, 3048F, 3049F, 3050F

## FOLLOW-UP CARE FOR CHILDREN PRESCRIBED ADHD MEDICATION (ADD)

<i>The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Two rates are reported:</i>		
<ul style="list-style-type: none"> <li><i>Initiation Phase. The percentage of members 6-12 years of age as of the Index Prescription Start Date (IPSD) with an ambulatory prescription dispensed for ADHD medication, who had one follow-up visit with practitioner with prescribing authority during the 30-day Initiation Phase.</i></li> <li><i>Continuation and Maintenance (C&amp;M) Phase. The percentage of members 6-12 years of age as of the IPSD with an ambulatory prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after Initiation Phase ended. One of the two visits (during the 31-300) may be a telephone visit with any practitioner.</i></li> </ul>		
CPT Code Set		
CPT codes	Stand Alone Visits: 96150-96154, 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99381-99384, 99391-99394, 99401-99404, 99411, 99412, 99510  Telephone Visits: 98966-98968, 99441-99443	
HCPCS	G0155, G0176, G0177, G0409-G0411, G0463, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, M0064, S0201, S9480, S9484, S9485, T1015	
CPT		POS
<i>Outpatient, intensive outpatient or partial hospitalization visits identified by the following CPT/POS codes must be with a practitioner with prescribing authority.</i>		
90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876	<i>WITH</i>	03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 71, 72
99221-99223, 99231-99233, 99238, 99239, 99251-99255	<i>WITH</i>	52, 53
ADHD Medications: <i>The following list of medications is based upon the 2017 HEDIS Tech Specs.</i>		
Description	Prescription	
CNS stimulants	<ul style="list-style-type: none"> <li>Amphetamine-dextroamphetamine</li> <li>Dexmethylphenidate</li> <li>Dextroamphetamine</li> <li>Lisdexamfetamine</li> <li>Methamphetamine</li> <li>Methylphenidate</li> </ul>	
Alpha-2 receptor agonists	<ul style="list-style-type: none"> <li>Clonidine</li> <li>Guanfacine</li> </ul>	
Miscellaneous ADHD medications	<ul style="list-style-type: none"> <li>Atomoxetine</li> </ul>	

## USE OF MULTIPLE CONCURRENT ANTIPSYCHOTICS IN CHILDREN AND ADOLESCENTS (APC)

The percentage of children and adolescents ages 1–17 who were on 2 or more concurrent antipsychotics in the measurement year. (Note: A lower rate indicates better performance.)

Antipsychotic Medications:				
Description	Prescription			
First Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>Chlorpromazine HCL</li> <li>Fluphenazine HCL</li> <li>Fluphenazine decanoate</li> <li>Haloperidol</li> </ul>	<ul style="list-style-type: none"> <li>Haloperidol decanoate</li> <li>Haloperidol lactate</li> <li>Loxapine HCL</li> </ul>	<ul style="list-style-type: none"> <li>Loxapine succinate</li> <li>Molindone HCL</li> <li>Perphenazine</li> <li>Pimozide</li> </ul>	<ul style="list-style-type: none"> <li>Thioridazine HCL</li> <li>Thiothixene</li> <li>Trifluoperazine HCL</li> </ul>
Second Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>Aripiprazole</li> <li>Asenapine</li> <li>Brexiprazole</li> <li>Cariprazine</li> </ul>	<ul style="list-style-type: none"> <li>Clozapine</li> <li>Iloperidone</li> <li>Lurasidone</li> <li>Olanzapine</li> </ul>	<ul style="list-style-type: none"> <li>Olanzapine pamoate</li> <li>Paliperidone</li> <li>Paliperidone palmitate</li> <li>Quetiapine fumarate</li> </ul>	<ul style="list-style-type: none"> <li>Risperidone</li> <li>Risperidone microspheres</li> <li>Ziprasidone HCL</li> <li>Ziprasidone mesylate</li> </ul>

## METABOLIC MONITORING FOR CHILDREN AND ADOLESCENTS ON ANTIPSYCHOTICS (APM)

The percentage of children and adolescents 1–17 years of age who had 2 or more antipsychotic prescriptions and had metabolic testing during the measurement year.

CPT Code Set				
Cholesterol Tests Other Than LDL	82465, 83718, 84478			
Glucose Tests	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951			
HbA1c Tests	83036, 83037, 3044F, 3045F, 3046F			
LDL-C Tests	80061, 83700, 83701, 83704, 83721, 3048F, 3049F, 3050F			
Antipsychotic Medications:				
Description	Prescription			
First Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>Chlorpromazine HCL</li> <li>Fluphenazine HCL</li> <li>Fluphenazine decanoate</li> <li>Haloperidol</li> </ul>	<ul style="list-style-type: none"> <li>Haloperidol decanoate</li> <li>Haloperidol lactate</li> <li>Loxapine HCL</li> </ul>	<ul style="list-style-type: none"> <li>Loxapine succinate</li> <li>Molindone HCL</li> <li>Perphenazine</li> <li>Pimozide</li> </ul>	<ul style="list-style-type: none"> <li>Thioridazine HCL</li> <li>Thiothixene</li> <li>Trifluoperazine HCL</li> </ul>
Second Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>Aripiprazole</li> <li>Asenapine</li> <li>Brexiprazole</li> <li>Cariprazine</li> </ul>	<ul style="list-style-type: none"> <li>Clozapine</li> <li>Iloperidone</li> <li>Lurasidone</li> <li>Olanzapine</li> </ul>	<ul style="list-style-type: none"> <li>Olanzapine pamoate</li> <li>Paliperidone</li> <li>Paliperidone palmitate</li> <li>Quetiapine fumarate</li> </ul>	<ul style="list-style-type: none"> <li>Risperidone</li> <li>Risperidone microspheres</li> <li>Ziprasidone HCL</li> <li>Ziprasidone mesylate</li> </ul>
Combinations	<ul style="list-style-type: none"> <li>Olanzapine-fluoxetine HCL (Symbyax)</li> <li>Perphenazine-amitriptyline HCL (Etrafon, Triavil [various])</li> </ul>			

## USE OF FIRST-LINE PSYCHOSOCIAL CARE FOR CHILDREN AND ADOLESCENTS ON ANTIPSYCHOTICS (APP)

The percentage of children and adolescents 1-17 years of age who had a new prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment.	
Psychosocial Care Code set	
ICD-9-CM Diagnosis	Schizophrenia: 295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95 Bipolar Disorder: 296.00-296.06, 296.10-296.16, 296.40-296.46, 296.50-296.56, 296.60-296.66, 296.7 Other Psychotic Disorders: 296.24, 296.34, 297.0, 297.1, 297.2, 297.3, 297.8, 297.9, 298.0, 298.1, 298.2, 298.3, 298.4, 298.8, 298.9, 299.00, 299.01, 299.10, 299.11, 299.80, 299.81, 299.90, 299.91, 307.20-307.23
ICD-10-CM Diagnosis	Schizophrenia: F20.0-F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Bipolar Disorder: F30.10-F30.13, F30.2-F30.4, F30.8, F30.9, F31.0, F31.10-F31.13, F31.2, F31.30-F31.32, F31.4, F31.5, F31.60-F31.64, F31.70-F31.78 Other Psychotic Disorders: F22, F23, F24, F28, F29, F32.3, F33.3, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F95.0, F95.1, F95.2, F95.8, F95.9
CPT Codes	Psychosocial Care: 90832-90834, 90836-90840, 90845-90847, 90849, 90853, 90875, 90876, 90880
HCPCS Codes	Psychosocial Care: G0176, G0177, G0409-G0411, H0004, H0035-H0040, H2000, H2001, H2011-H2014, H2017-H2020, S0201, S9480, S9484, S9485
Antipsychotic Medications:	
Description	Prescription
First Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>• Chlorpromazine HCL</li> <li>• Fluphenazine HCL</li> <li>• Fluphenazine decanoate</li> <li>• Haloperidol</li> <li>• Haloperidol decanoate</li> <li>• Haloperidol lactate</li> <li>• Loxapine HCL</li> <li>• Loxapine succinate</li> <li>• Moliindone HCL</li> <li>• Perphenazine</li> <li>• Pimozide</li> <li>• Thioridazine HCL</li> <li>• Thiothixene</li> <li>• Trifluoperazine HCL</li> </ul>
Second Generation Antipsychotic Medications	<ul style="list-style-type: none"> <li>• Aripiprazole</li> <li>• Asenapine</li> <li>• Brexpiprazole</li> <li>• Cariprazine</li> <li>• Clozapine</li> <li>• Iloperidone</li> <li>• Lurasidone</li> <li>• Olanzapine</li> <li>• Olanzapine pamoate</li> <li>• Paliperidone</li> <li>• Paliperidone palmitate</li> <li>• Quetiapine fumarate</li> <li>• Risperidone</li> <li>• Risperidone microspheres</li> <li>• Ziprasidone HCL</li> <li>• Ziprasidone mesylate</li> </ul>
Combinations	<ul style="list-style-type: none"> <li>• Olanzapine-fluoxetine HCL (Symbyax)</li> <li>• Perphenazine-amitriptyline HCL (Etrafon, Triavil [various])</li> </ul>

## FOLLOW-UP AFTER EMERGENCY DEPARTMENT VISIT FOR MENTAL ILLNESS (FUM)

<b>FIRST-YEAR MEASURE</b>	
<p>The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness, who had a follow-up visit for mental illness. Two rates are reported:</p> <ol style="list-style-type: none"> <li>The percentage of ED visits for which the member received follow-up within 30 days of the ED visit.</li> <li>The percentage of ED visits for which the member received follow-up within 7 days of the ED visit.</li> </ol> <p>The follow-up visit after the ED visit can be with any practitioner.</p>	
ICD-9-CM Diagnosis	290.0, 290.10-290.13, 290.20, 290.21, 290.3, 290.40-290.43, 290.8, 290.9, 293.0, 293.1, 293.81-293.84, 293.89, 293.9, 294.0, 294.10, 294.11, 294.20, 294.21, 294.8, 294.9, 295.00, 295.01-295.05, 295.10-295.15, 295.20-295.25, 295.30-295.35, 295.40-295.45, 295.50-295.55, 295.60-295.65, 295.70-295.75, 295.80-295.85, 295.90-295.95, 296.00-296.06, 296.10-296.16, 296.20-296.26, 296.30-296.36, 296.40-296.46, 296.50-296.56, 296.60-296.66, 296.7, 296.80-296.82, 296.89, 296.90, 296.99, 297.0, 297.1, 297.2, 297.3, 297.8, 297.9, 298.0, 298.1, 298.2, 298.3, 298.4, 298.8, 298.9, 299.00, 299.01, 299.10, 299.11, 299.80, 299.81, 299.90, 299.91, 300.00-300.02, 300.09-300.16, 300.19-300.23, 300.29, 300.3, 300.4, 300.5, 300.6, 300.7, 300.81, 300.82, 300.89, 300.9, 301.0, 301.10-301.13, 301.20-301.22, 301.3, 301.4, 301.5, 301.51, 301.59, 301.6, 301.7, 301.81-301.84, 301.89, 301.9, 302.0, 302.1, 302.2, 302.3, 302.4, 302.50-302.53, 302.6, 302.70-302.76, 302.79, 302.81-302.85, 302.89, 302.9, 306.0, 306.1, 306.2, 306.3, 306.4, 306.4, 306.50-306.53, 306.59, 306.6, 306.7, 306.8, 306.9, 307.0, 307.1, 307.20-307.23, 307.3, 307.40-307.54, 307.59, 307.6, 307.7, 307.80, 307.81, 307.89, 307.9, 308.0, 308.1, 308.2, 308.3, 308.4, 308.9, 309.1, 309.21-309.24, 309.28, 309.29, 309.3, 309.4, 309.81-309.83, 309.89, 309.9, 310.0, 310.1, 310.2, 310.81, 310.89, 310.9, 311, 312.00-312.03, 312.10-312.13, 312.20-312.23, 312.30-312.35, 312.39, 312.4, 312.81, 312.82, 312.89, 312.9, 313.0, 313.1, 313.21-313.23, 313.3, 313.81-313.83, 313.89, 313.9, 314.00, 314.01, 314.1, 314.2, 314.8, 314.9, 315.00-315.02, 315.09, 315.1, 315.2, 315.31-315.35, 315.39, 315.4, 315.5, 315.8, 315.9, 316
ICD-10-CM Diagnosis	Use the appropriate code family: F
CPT	HCPS
Follow-up visits identified by the following CPT or HCPCS codes with any practitioner within 7 and 30 days of ED Visit.	
Stand Alone Visits: 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99383-99387, 99393-99397, 99401-99404, 99411, 99412, 99495, 99496, 99510 Telehealth: 98966-98969, 99441-99444	Stand Alone Visits: G0155, G0176, G0177, G0409-G0411, G0463, H0002, H0004, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, M0064, S0201, S9480, S9484, S9485, T1015
CPT	POS
Follow-up visits identified by the following CPT with POS codes with any practitioner within 7 and 30 days of ED Visit.	
Follow-Up Group 1: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90867-90870, 90875, 90876	WITH 03, 05, 07, 09, 11-20, 22, 24, 33, 49, 50, 52, 53, 71, 72
Follow-Up Group 2: 99221-99223, 99231-99233, 99238, 99239, 99251-99255	WITH 52, 53

## FOLLOW-UP AFTER EMERGENCY DEPARTMENT VISIT FOR ALCOHOL AND OTHER DRUG DEPENDENCE (FUA)

<b>FIRST-YEAR MEASURE</b>	
<p>The percentage of emergency department (ED) visits for members 13 years of age and older with a principal diagnosis of alcohol or other drug (AOD) dependence, who had a follow up visit for AOD. Two rates are reported:</p> <ol style="list-style-type: none"> <li>The percentage of ED visits for which the member received follow-up within 30 days of the ED visit.</li> <li>The percentage of ED visits for which the member received follow-up within 7 days of the ED visit.</li> </ol> <p>The follow-up visit after the ED visit can be with any practitioner.</p>	
ICD-9-CM Diagnosis	AOD: 291.0-291.5, 291.81, 291.82, 291.89, 291.9, 303.00-303.02, 303.90-303.92, 304.00-304.02, 304.10-304.12, 304.20-304.22, 304.30-304.32, 304.40-304.42, 304.50-304.52, 304.60-304.62, 304.70-304.72, 304.80-304.82, 304.90-304.92, 305.00-305.02, 305.20-305.22, 305.30-305.32, 305.40-305.42, 305.50-305.52, 305.60-305.62, 305.70-305.72, 305.80-305.82, 305.90-305.92, 535.30, 535.31, 571.1
ICD-10-CM Diagnosis	Use the appropriate code family: F
CPT	HCPS
Follow-up visits identified by the following CPT or HCPCS codes with any practitioner within 7 and 30 days of ED Visit.	
Stand Alone Visits: 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99384-99387, 99394-99397, 99401-99404, 99408, 99409, 99411, 99412, 99510 Telehealth: 98966-98969, 99441-99444	Stand Alone Visits: G0155, G0176, G0177, G0396, G0397, G0409-G0411, G0443, G0463, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0020, H0022, H0031, H0034-H0037, H0039, H0040, H2000, H2001, H2010-H2020, H2035, H2036, M0064, S0201, S9480, S9484, S9485, T1006, T1012, T1015
CPT	POS
Follow-up visits identified by the following CPT with POS codes with any practitioner within 7 and 30 days of ED Visit.	



Follow-Up Group 1: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876	<i>WITH</i>	03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 71, 72
Follow-Up Group 2: 99221-99223, 99231-99233, 99238, 99239, 99251-99255	<i>WITH</i>	52, 53